education

DEPARTMENT: EDUCATION
MPUMALANGA PROVINCE

NATIONAL SENIOR CERTIFICATE

GRADE 12

PHYSICAL SCIENCES: CHEMISTRY (P2)

SEPTEMBER 2016

MARKS: 150

TIME: 3 hours

This question paper consists of 18 pages and 4 data sheets.

INSTRUCTIONS AND INFORMATION

1. Write your name in the appropriate space on the ANSWER BOOK.
2. This question paper consists of TEN questions. Answer ALL the questions in the ANSWER BOOK.
3. Start EACH question on a NEW page in the ANSWER BOOK.
4. Number the answers correctly according to the numbering system used in this question paper.
5. Leave ONE line between two sub questions, for example between QUESTION 2.1 and QUESTION 2.2.
6. You may use a non-programmable calculator.
7. You may use appropriate mathematical instruments.
8. You are advised to use the attached DATA SHEETS.
9. Show ALL formulae and substitutions in ALL calculations.
10. Round off your final numerical answers to a MINIMUM of TWO decimal places.
11. Give brief motivations, discussions, et cetera where required.
12. Write neatly and legibly.

QUESTION 1 : MULTIPLE-CHOICE QUESTIONS

Four options are provided as possible answers to the following questions. Each question has only ONE correct answer. Choose the answer and write only the letter (A – D) next to the question number (1.1 – 1.10) in the ANSWER BOOK , for example, 1.11 E.

1.1 Which ONE of the following elements is a non-mineral nutrient?

A Mg

B Na

C C

D Zn (2)

1.2 Which ONE of the following is a catalyst used in the Haber process?

A Silica gel

B Iron oxide

C Vanadium pentoxide

D Nickel (2)

1.3 Consider the following reaction in equilibrium in a closed container.

$$4\ NH_3\ (g) + 3\ O_2\ (g) \rightleftharpoons 6\ H_2O\ (g) + 2\ N_2(g) \qquad \Delta H < 0$$

Which ONE of the following changes in the conditions at equilibrium will result in an increase in the value of the equilibrium constant, K_c?

A Increasing the ammonia concentration

B Increasing the temperature

C Decreasing the temperature

D Adding a platinum catalyst (2)

1.4 Which type of reaction is illustrated by the following chemical equation

$$CH_2CHCH_3 + H_2 \rightarrow CH_3CH_2CH_3$$

A Esterification

B Addition

C Elimination

D Substitution (2)

1.5 A compound with molecular formula $C_4H_8O_2$ could be

(i) an alcohol

(ii) a carboxylic acid

(iii) an ester

Which ONE of the following statements is CORRECT?

A (i) only

B (i) and (ii) only

C (i) and (iii) only

D (ii) and (iii) only (2)

1.6 Ammonium sulphate ($(NH_4)_2SO_4$) is dissolved in water. Which ONE of the following statements regarding the solution formed, is CORRECT?

A pH = 7

B $[H_3O^+] \cdot [OH^-] < 1 \times 10^{-14}$

C $[H_3O^+] > [OH^-]$

D $[H_3O^+] < [OH^-]$ (2)

1.7 Consider the reversible reaction represented below:

$$CH_3COOH(aq) + HSO_4^-(aq) \rightleftharpoons CH_3COO^-(aq) + H_2SO_4(aq)$$

The strongest base in the reaction above is:

A CH_3COO^-

B CH_3COOH

C HSO_4^-

D H_2SO_4 (2)

1.8 Which ONE of the following will be changed by the addition of a catalyst?

A The amount of products present at equilibrium

B The concentration of reactants present at equilibrium

C The length of time for which the reaction remains at equilibrium

D The time it takes to reach equilibrium (2)

1.9 A hypothetical electrochemical cell is constructed by combining two half-cells. The two half-reactions and electrode potentials are given below:

$X(s) + 2e^- \rightleftharpoons X^{2-}(aq)$ $E^\theta = +1,5\ V$
$Y^+(aq) + e^- \rightleftharpoons Y(s)$ $E^\theta = -1,5\ V$

Which ONE of the following statements is TRUE?

A $X^{2-}(aq)$ will be oxidised more readily than Y(s)

B $Y^+(aq)$ will be one of the products formed in the cell reaction

C The mass of X will increase when the cell is delivering current

D Electrons will flow from X (s) to Y (s) in the external circuit when the cell is delivering current (2)

1.10 A Zn – Cu cell has been operating for some time but the net cell reaction has not reached equilibrium. How will an increase in the $[Cu^{2+}]$ affect the emf of the cell?

A The emf will increase

B The emf will decrease






C The emf will remain the same

D The emf will be zero (2)

[20]

QUESTION 2 (Start on a new page)

Six organic compounds (**A – F**) are given in the table below.

A		**B**	
C		**D**	Pentanoic acid
E		**F**	

2.1 Write down the:

2.1.1 Letter of the compound that represents an aldehyde (1)

2.1.2 Structural formula of the functional group of compound **D** (1)

2.1.3 IUPAC name of compound **C** (2)

2.1.4 Letter of the compound that represents a tertiary alcohol (1)

2.1.5 Letters of compounds that represents TWO functional isomers (2)

2.1.6 To which homologous series does compound **B** belong? (1)

2.1.7 Is compound **B** SATURATED or UNSATURATED? Explain the answer. (3)

2.2 Compound **F** is prepared in the laboratory.

2.2.1 How can one quickly establish whether compound **F** is indeed being formed without performing a chemical test? (1)

2.2.2 Give the IUPAC name of the alcohol needed to prepare compound **F**. (2)

[14]

QUESTION 3 (Start on a new page)

The boiling points of pentane, 2-methylbutane and 2,2-dimethylpropane were determined during a practical investigation and recorded in the table below.

ALKANE	MOLECULAR MASS ($g \cdot mol^{-1}$)	BOILING POINT (°C)
Pentane	72	36
2-methylbutane	72	28
2,2-dimethylpropane	72	10

3.1 Define the term *boiling point.* (2)

3.2 State a suitable hypothesis for the investigation. (2)

3.3 Fully explain why pentane has the highest boiling point. (4)


3.4 Which ONE of the three compounds in the table has the highest vapour pressure? Give a reason for the answer (2)

3.5 Write down the balanced equation for the complete combustion of pentane. (3)

[13]

QUESTION 4 (Start on a new page)

4.1 The flow diagram shows how two compounds can be formed from 2-bromopentane.



Reaction 1 takes place in the presence of concentrated sodium hydroxide.
Write down:

4.1.1 An additional reaction condition for this reaction (1)

4.1.2 The type of reaction of which **Reaction 1** is an example (1)

4.1.3 The **structural formula** of compound A (2)

Reaction 2 takes place in the presence of a certain inorganic compound.
Write down the:

4.1.4 NAME of the inorganic compound (1)

4.1.5 Type of reaction of which **Reaction** 2 is an example. (1)

Reaction 3 takes place in the presence of concentrated sulphuric acid and heat.

4.1.6 Write down the type of reaction that converts pentan-2-ol to compound A. (1)

4.2 A learner is preparing an ester using ethanol (molecular formula: C_2H_6O) and methanoic acid (molecular formula: CH_2O_2).

The balanced equation for the reaction is given below:


$$C_2H_6O + CH_2O_2 \rightarrow C_3H_6O_2 + H_2O$$

4.2.1 What type of reaction is represented in the equation above? (1)

4.2.2 Write down the NAME or FORMULA of the catalyst used in this reaction. (1)

4.2.3 When 60 g of **impure** ethanol fully reacts with excess methanoic acid, 68 g of $C_3H_6O_2$ is produced.
Calculate the percentage purity of ethanol. (6)

4.3 The structural formula of a polymer is shown below:



Write down the:

4.3.1 STRUCTURAL FORMULA of the monomer that is used to prepare the above polymer. (2)

4.3.2 Type of polymerisation reaction (ADDITION or CONDENSATION) that is used to prepare this polymer. (1)

[18]

QUESTION 5 (Start on a new page)

A group of leaners use the reaction between zinc and sulphuric acid to investigate one of the **factors** that affects reaction rate.

They add 6,5 g of zinc granules to excess **DILUTE** sulphuric acid and measure the mass of zinc used per unit time.

The learners then repeat the experiment using the same mass of zinc granules with excess **CONCENTRATED** sulphuric acid.


The balanced equation for the reaction is:

$$Zn(s) + H_2SO_4(aq) \rightarrow ZnSO_4(aq) + H_2(g)$$

5.1 Define the term *reaction rate* (in words). (2)

5.2 Write down the independent variable. (1)

5.3 The results obtained for the reaction using DILUTE sulphuric acid are represented in the graph below:




Use the graph to answer the questions below:

5.3.1 Calculate the average rate of the reaction during the first 60 s in **gram per second** (5)

5.3.2 Compare the rate of reaction at time = 140 s to the rate of reaction at time = 60 s.
Choose from GREATER THAN, LESS THAN or EQUAL TO. (1)

The Maxwell –Boltzmann distribution curve below represents the number of particles against kinetic energy for the reaction of zinc with **DILUTE** sulphuric acid. The graph is labelled **A**.



5.4 Redraw the curve in the ANSWER BOOK. On the same set of axes, sketch the curve that will be obtained for the reaction with **CONCENTRATED** sulphuric acid. Clearly label the curves **A** and **B**. (2)

5.5. Use the collision theory to explain how an increase in the **concentration** of sulphuric acid will affect the rate of reaction (3)

[14]

QUESTION 6 (Start on a new page)

2 moles of carbon dioxide (CO_2), 1 mole of hydrogen (H_2) and 0,2 mole of carbon monoxide (CO) are sealed in a 2 dm^3 container. The reaction reaches equilibrium at 1 000°C. The balanced equation for the reaction is:

$$CO_2(g) + H_2(g) \rightleftharpoons CO(g) + H_2O(g)$$

On analysis of the equilibrium mixture , it is found that the concentration of the water vapour is 0,362 $mol.dm^{-3}$.

6.1 Calculate the equilibrium constant (Kc) at 1 000°C for this reaction. (7)

6.2 It is found that at a temperature of 1 500°C, the value of the equilibrium constant calculated in QUESTION 6.1, decreases. Is the forward reaction EXOTHERMIC or ENDOTHERMIC? Fully explain how you arrived at the answer. (4)

6.3 An additional 0,277 mole of water vapour is pumped into the sealed 2 dm^3 container. The volume of the container stays the same. Equillibrium is reinstated at 1 000°C.

6.3.1 How is the concentration of the CO (g) affected?
Choose from INCREASES, DECREASES or STAYS THE SAME. (1)

6.3.2 Use Le Chatelier's Principle to explain the answer in QUESTION 6.3.1. (3)

[15]

QUESTION 7 (Start on a new page)

7.1 Sulphuric acid is a **strong** diprotic acid.

Define the term *strong acid* (2)

7.2 HSO_4^- can behave either as an acid or a base.

7.2.1 Give the term that is used for substances such as HSO_4^-. (1)

7.2.2 Write down the balanced equation for the reaction of HSO_4^- with H_2O in which HSO_4^- acts as an acid. (2)

7.3 Calculate the pH of a $Mg(OH)_2$ solution with a concentration of 0,2 $mol{\cdot}dm^{-3}$. Assume that $Mg(OH)_2$ dissociates completely. (4)

7.4 Vinegar is (3 – 5)% ethanoic acid by mass, with the remaining mass being a solvent. To determine the percentage of ethanoic acid in a vinegar sample, it is titrated against a NaOH solution. The balanced equation for the reaction is:

$$CH_3COOH(aq) + NaOH(aq) \rightarrow CH_3COONa(aq) + H_20(\ell)$$

A 10 g sample of vinegar is titrated against a 0,309 $mol{\cdot}dm^{-3}$ NaOH solution. 19,57 cm^3 of NaOH is required to reach end point.

7.4.1 Determine the mass of the ethanoic acid in the 10 g sample. (5)

7.4.2 Show by calculation that the percentage of the ethanoic acid is within the given range. (2)

The table below provides information of three different indicators:

INDICATOR	COLOUR CHANGE	COLOUR CHANGE pH RANGE
Methyl orange	Red – yellow	3,0 – 4,4
Bromothymol blue	Yellow – blue	6,0 – 7,6
Phenolphthalein	Colourless - pink	8,2 – 9,8

7.4.3 Use the information in the table above and choose a suitable indicator to use in the above titration. Give a reason for the answer. (2)

7.4.4 The ethanoate ions that form during the reaction, react with water according to the balanced equation below:


$$CH_3COO^-(aq) + H_2O(\ell) \rightleftharpoons CH_3COOH(aq) + OH^-(aq)$$

Write down the name of the process described by the underlined phrase. (1)

[19]

QUESTION 8 (Start on a new page)

The two half-cells, Fe(s) | Fe^{2+}(aq) and O_2(g) | $H_2O(\ell)$ in acid solution, are used to set up an electrochemical cell. The cell operates under standard conditions.



8.1 Write down TWO standard conditions, other than temperature, which apply to this electrochemical cell. (2)

8.2 Give TWO reasons, besides being a solid, why platinum is a suitable electrode in the above cell. (2)

8.3 Write down the half-reaction that takes place at the cathode of this cell. (2)

8.4 Write down the cell notation of this cell. (3)


8.5 Calculate the initial emf of this cell. (4)

8.6 The reading of the voltmeter becomes zero after using the cell for several hours. Give a reason for this reading by referring to the cell reaction. (1)

[14]

QUESTION 9 (start on a new page)

The diagram below show an electrolytic cell used to coat a nickel coin with silver.



9.1 Which electrode (anode or cathode) does the nickel coin represent? (1)

9.2 Define the term *electrolyte*. (2)

9.3 Write down the NAME or FORMULA of a suitable electrolyte used in this cell. (1)

9.4 Write down the half-reaction that takes place on the surface of the nickel coin. (2)


9.5 How will the concentration of the electrolyte change during electrolysis? Choose from INCREASE, DECREASE or REMAINS THE SAME. (1)

9.6 Explain the answer in QUESTION 9.5. (2)

[9]

QUESTION 10 (Start on a new page)

The diagram below illustrates the processes used to manufacture a certain fertilizer. Study the diagram and answer the questions that follow.



10.1.1 NAME or FORMULA of substance A that is obtained by the fractional distillation of liquid air. (1)

10.1.2 Balanced equation for the reaction that occurs in **Process 1.** (3)


10.1.3 NAME of the substance B that forms as a result of **Process 1.** (1)

10.1.4 NAME of **process 2.** (1)

10.1.5 NAME or FORMULA of fertiliser **C.** (1)

10.1.6 NAME of the process that occurs when excess fertiliser **C** runs into the rivers. (1)

10.2 A farmer has two bags of fertiliser marked **A** and **B**, as shown below.



10.2.1 The farmer wants to grow tomatoes and fruits. Which ONE of these fertilisers must the farmer use, write down **A** or **B**.
Give a reason for the answer. (2)

10.2.2 Calculate the mass of potassium in bag **B.** (4)

[14]

TOTAL MARKS: 150

DATA FOR PHYSICAL SCIENCES GRADE 12
PAPER 2 (CHEMISTRY)

GEGEWENS VIR FISIESE WETENSKAPPE GRAAD 12
VRAESTEL 2 (CHEMIE)

TABLE 1: PHYSICAL CONSTANTS/*TABEL 1: FISIESE KONSTANTES*

NAME/*NAAM*	SYMBOL/*SIMBOOL*	VALUE/*WAARDE*
Standard pressure *Standaarddruk*	p^θ	$1,013 \times 10^5$ Pa
Molar gas volume at STP *Molêre gasvolume by STD*	V_m	$22,4\ dm^3 \cdot mol^{-1}$
Standard temperature *Standaardtemperatuur*	T^θ	273 K
Charge on electron *Lading op elektron*	e	$-1,6 \times 10^{-19}$ C
Avogadro's constant *Avogadro-konstante*	N_A	$6,02 \times 10^{23}\ mol^{-1}$

TABLE 2: FORMULAE/*TABEL 2: FORMULES*

$n = \frac{m}{M}$	$n = \frac{N}{N_A}$
$c = \frac{n}{V}$ or/*of* $c = \frac{m}{MV}$	$n = \frac{V}{V_m}$
$\frac{c_a V_a}{c_b V_b} = \frac{n_a}{n_b}$	$pH = -\log[H_3O^+]$
$K_w = [H_3O^+][OH^-] = 1 \times 10^{-14}$ at/*by* 298 K	
$E^\theta_{cell} = E^\theta_{cathode} - E^\theta_{anode}$ / $E^\theta_{sel} = E^\theta_{katode} - E^\theta_{anode}$ or/*of* $E^\theta_{cell} = E^\theta_{reduction} - E^\theta_{oxidation}$ / $E^\theta_{sel} = E^\theta_{reduksie} - E^\theta_{oksidasie}$ or/*of* $E^\theta_{cell} = E^\theta_{oxidising\ agent} - E^\theta_{reducing\ agent}$ / $E^\theta_{sel} = E^\theta_{oksideermiddel} - E^\theta_{reduseermiddel}$	

TABLE 3: THE PERIODIC TABLE OF ELEMENTS
TABEL 3: DIE PERIODIEKE TABEL VAN ELEMENTE




TABLE 4A: STANDARD REDUCTION POTENTIALS
TABEL 4A: STANDAARD-REDUKSIEPOTENSIALE

Half-reactions/*Halfreaksies*			$E^{\ominus}$ (V)
$F_2(g) + 2e^-$	⇌	$2F^-$	+ 2,87
$Co^{3+} + e^-$	⇌	Co^{2+}	+ 1,81
$H_2O_2 + 2H^+ + 2e^-$	⇌	$2H_2O$	+ 1,77
$MnO_4^- + 8H^+ + 5e^-$	⇌	$Mn^{2+} + 4H_2O$	+ 1,51
$Cl_2(g) + 2e^-$	⇌	$2Cl^-$	+ 1,36
$Cr_2O_7^{2-} + 14H^+ + 6e^-$	⇌	$2Cr^{3+} + 7H_2O$	+ 1,33
$O_2(g) + 4H^+ + 4e^-$	⇌	$2H_2O$	+ 1,23
$MnO_2 + 4H^+ + 2e^-$	⇌	$Mn^{2+} + 2H_2O$	+ 1,23
$Pt^{2+} + 2e^-$	⇌	Pt	+ 1,20
$Br_2(l) + 2e^-$	⇌	$2Br^-$	+ 1,07
$NO_3^- + 4H^+ + 3e^-$	⇌	$NO(g) + 2H_2O$	+ 0,96
$Hg^{2+} + 2e^-$	⇌	Hg(l)	+ 0,85
$Ag^+ + e^-$	⇌	Ag	+ 0,80
$NO_3^- + 2H^+ + e^-$	⇌	$NO_2(g) + H_2O$	+ 0,80
$Fe^{3+} + e^-$	⇌	Fe^{2+}	+ 0,77
$O_2(g) + 2H^+ + 2e^-$	⇌	H_2O_2	+ 0,68
$I_2 + 2e^-$	⇌	$2I^-$	+ 0,54
$Cu^+ + e^-$	⇌	Cu	+ 0,52
$SO_4^{2-} + 4H^+ + 4e^-$	⇌	$S + 2H_2O$	+ 0,45
$2H_2O + O_2 + 4e^-$	⇌	$4OH^-$	+ 0,40
$Cu^{2+} + 2e^-$	⇌	Cu	+ 0,34
$SO_4^{2-} + 4H^+ + 2e^-$	⇌	$SO_2(g) + 2H_2O$	+ 0,17
$Cu^{2+} + e^-$	⇌	Cu^+	+ 0,16
$Sn^{4+} + 2e^-$	⇌	Sn^{2+}	+ 0,15
$S + 2H^+ + 2e^-$	⇌	$H_2S(g)$	+ 0,14
$2H^+ + 2e^-$	⇌	$H_2(g)$	0,00
$Fe^{3+} + 3e^-$	⇌	Fe	− 0,06
$Pb^{2+} + 2e^-$	⇌	Pb	− 0,13
$Sn^{2+} + 2e^-$	⇌	Sn	− 0,14
$Ni^{2+} + 2e^-$	⇌	Ni	− 0,27
$Co^{2+} + 2e^-$	⇌	Co	− 0,28
$Cd^{2+} + 2e^-$	⇌	Cd	− 0,40
$Cr^{3+} + e^-$	⇌	Cr^{2+}	− 0,41
$Fe^{2+} + 2e^-$	⇌	Fe	− 0,44
$Cr^{3+} + 3e^-$	⇌	Cr	− 0,74
$Zn^{2+} + 2e^-$	⇌	Zn	− 0,76
$2H_2O + 2e^-$	⇌	$H_2(g) + 2OH^-$	− 0,83
$Cr^{2+} + 2e^-$	⇌	Cr	− 0,91
$Mn^{2+} + 2e^-$	⇌	Mn	− 1,18
$Al^{3+} + 3e^-$	⇌	Al	− 1,66
$Mg^{2+} + 2e^-$	⇌	Mg	− 2,36
$Na^+ + e^-$	⇌	Na	− 2,71
$Ca^{2+} + 2e^-$	⇌	Ca	− 2,87
$Sr^{2+} + 2e^-$	⇌	Sr	− 2,89
$Ba^{2+} + 2e^-$	⇌	Ba	− 2,90
$Cs^+ + e^-$	⇌	Cs	− 2,92
$K^+ + e^-$	⇌	K	− 2,93
$Li^+ + e^-$	⇌	Li	− 3,05

Increasing oxidising ability/*Toenemende oksiderende vermoë* ↑

Increasing reducing ability/*Toenemende reduserende vermoë* ↓

TABLE 4B: STANDARD REDUCTION POTENTIALS
TABEL 4B: STANDAARD-REDUKSIEPOTENSIALE

Half-reactions/*Halfreaksies*			$E^{\ominus}$ (V)
$Li^+ + e^-$	⇌	Li	− 3,05
$K^+ + e^-$	⇌	K	− 2,93
$Cs^+ + e^-$	⇌	Cs	− 2,92
$Ba^{2+} + 2e^-$	⇌	Ba	− 2,90
$Sr^{2+} + 2e^-$	⇌	Sr	− 2,89
$Ca^{2+} + 2e^-$	⇌	Ca	− 2,87
$Na^+ + e^-$	⇌	Na	− 2,71
$Mg^{2+} + 2e^-$	⇌	Mg	− 2,36
$Al^{3+} + 3e^-$	⇌	Al	− 1,66
$Mn^{2+} + 2e^-$	⇌	Mn	− 1,18
$Cr^{2+} + 2e^-$	⇌	Cr	− 0,91
$2H_2O + 2e^-$	⇌	$H_2(g) + 2OH^-$	− 0,83
$Zn^{2+} + 2e^-$	⇌	Zn	− 0,76
$Cr^{3+} + 3e^-$	⇌	Cr	− 0,74
$Fe^{2+} + 2e^-$	⇌	Fe	− 0,44
$Cr^{3+} + e^-$	⇌	Cr^{2+}	− 0,41
$Cd^{2+} + 2e^-$	⇌	Cd	− 0,40
$Co^{2+} + 2e^-$	⇌	Co	− 0,28
$Ni^{2+} + 2e^-$	⇌	Ni	− 0,27
$Sn^{2+} + 2e^-$	⇌	Sn	− 0,14
$Pb^{2+} + 2e^-$	⇌	Pb	− 0,13
$Fe^{3+} + 3e^-$	⇌	Fe	− 0,06
$2H^+ + 2e^-$	⇌	$H_2(g)$	0,00
$S + 2H^+ + 2e^-$	⇌	$H_2S(g)$	+ 0,14
$Sn^{4+} + 2e^-$	⇌	Sn^{2+}	+ 0,15
$Cu^{2+} + e^-$	⇌	Cu^+	+ 0,16
$SO_4^{2-} + 4H^+ + 2e^-$	⇌	$SO_2(g) + 2H_2O$	+ 0,17
$Cu^{2+} + 2e^-$	⇌	Cu	+ 0,34
$2H_2O + O_2 + 4e^-$	⇌	$4OH^-$	+ 0,40
$SO_2 + 4H^+ + 4e^-$	⇌	$S + 2H_2O$	+ 0,45
$Cu^+ + e^-$	⇌	Cu	+ 0,52
$I_2 + 2e^-$	⇌	$2I^-$	+ 0,54
$O_2(g) + 2H^+ + 2e^-$	⇌	H_2O_2	+ 0,68
$Fe^{3+} + e^-$	⇌	Fe^{2+}	+ 0,77
$NO_3^- + 2H^+ + e^-$	⇌	$NO_2(g) + H_2O$	+ 0,80
$Ag^+ + e^-$	⇌	Ag	+ 0,80
$Hg^{2+} + 2e^-$	⇌	Hg(l)	+ 0,85
$NO_3^- + 4H^+ + 3e^-$	⇌	$NO(g) + 2H_2O$	+ 0,96
$Br_2(l) + 2e^-$	⇌	$2Br^-$	+ 1,07
$Pt^{2+} + 2e^-$	⇌	Pt	+ 1,20
$MnO_2 + 4H^+ + 2e^-$	⇌	$Mn^{2+} + 2H_2O$	+ 1,23
$O_2(g) + 4H^+ + 4e^-$	⇌	$2H_2O$	+ 1,23
$Cr_2O_7^{2-} + 14H^+ + 6e^-$	⇌	$2Cr^{3+} + 7H_2O$	+ 1,33
$Cl_2(g) + 2e^-$	⇌	$2Cl^-$	+ 1,36
$MnO_4^- + 8H^+ + 5e^-$	⇌	$Mn^{2+} + 4H_2O$	+ 1,51
$H_2O_2 + 2H^+ + 2e^-$	⇌	$2H_2O$	+ 1,77
$Co^{3+} + e^-$	⇌	Co^{2+}	+ 1,81
$F_2(g) + 2e^-$	⇌	$2F^-$	+ 2,87

Increasing oxidising ability/*Toenemende oksiderende vermoë* ↓

Increasing reducing ability/*Toenemende reduserende vermoë* ↑