education

Department:
Education
PROVINCE OF KWAZULU-NATAL

NATIONAL SENIOR CERTIFICATE

GRADE 12

PHYSICAL SCIENCES P2 (CHEMISTRY)

PREPARATORY EXAMINATION

SEPTEMBER 2019

MARKS : 150

TIME : 3 Hours

This question paper consists of 14 pages, a special answer sheet and 4 data sheets.

Copyright reserved

INSTRUCTIONS AND INFORMATION

1. Write your centre number and examination number in the appropriate spaces on the ANSWER BOOK.
2. This question paper consists of TEN questions. Answer ALL the questions in the ANSWER BOOK.
3. Start EACH question on a NEW page in the ANSWER BOOK.
4. Number the answers correctly according to the numbering system used in this question paper.
5. Leave ONE line between two subquestions, for example between QUESTION 2.1 and QUESTION 2.2.
6. You may use a non-programmable calculator.
7. You may use appropriate mathematical instruments.
8. You are advised to use the attached DATA SHEETS.
9. Show ALL formulae and substitutions in ALL calculations.
10. Round off your final numerical answers to a minimum of TWO decimal places.
11. Give brief motivations, discussions et cetera where required.
12. Write neatly and legibly.

Copyright reserved

QUESTION 1: MULTIPLE-CHOICE QUESTIONS

Various options are provided as possible answers to the following questions. Choose the answer and write only the letter (A-D) next to the question number (1.1-1.10) in the ANSWER BOOK, for example 1.11 D.

1.1 Which ONE of the following organic reactions below involves the bonding of monomers by a dehydration process?

A Halogenation.

B Dehydropolymerisation.

C Addition polymerisation.

D Condensation polymerisation. (2)

1.2 A compound with the molecular formula $C_5H_{10}O$ could be . . .

(I) an ester.
(II) a ketone.
(III) an aldehyde.

A (I) only.

B (I) and (II) only

C (II) and (III) only

D (I), (II) and (III) (2)

1.3 A catalyst will change the . . .

A time required to reach equilibrium.

B amount of products present at equilibrium.

C amount of reactants present at equilibrium.

D length time for a reaction remains at equilibrium. (2)

1.4 The difference between the potential energy of the products and the potential energy of the reactants in a chemical reaction is equal to the . . .

A rate of the reaction.

B enthalpy of the reaction.

C enthalpy change of the reaction.

D total potential energy of the particles. (2)

Copyright reserved

1.5 Which ONE of the following represents the products formed during the hydrolysis of sodium ethanoate, $NaCH_3COO$?

A $Na^+(aq)$ and $OH^-(aq)$

B $Na^+(aq)$ and $CH_3COO^-(aq)$

C $H_3O^+(aq)$ and $CH_3COO^-(aq)$

D $CH_3COOH(aq)$ and $OH^-(aq)$ (2)

1.6 The reaction, $H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$, $\Delta H < 0$, reaches equilibrium at a temperature of 445 °C in a sealed container.

Which ONE of the following change/s will INCREASE the CONCENTRATION of HI(g)?

I adding a catalyst
II decreasing the temperature
III increasing the pressure by reducing the volume of the reaction container

A only II


B only III

C I and II only

D II and III only (2)

Copyright reserved

1.7 Learners set up an electrochemical cell, shown in the simplified diagram below, using magnesium and lead electrodes.



Which of the following changes will INCREASE the reading on the voltmeter?

(i) Decreasing the initial concentration of the electrolyte in half-cell A.

(ii) Decreasing the initial concentration of the electrolyte in half-cell B.

(iii) Increasing the initial concentration of the electrolyte in half-cell A.

(iv) Increasing the initial concentration of the electrolyte in half-cell B.

A (i) Only

B (ii) only

C (ii) and (iii) only

D (i) and (iv) only (2)

1.8 The balanced equation for the net(overall) cell reaction that occurs during the electrolysis of a concentrated sodium chloride solution is :

$$2\ H_2O(\ell) + 2\ Cl^- \rightarrow Cl_2(g) + H_2(g) + 2OH^-(aq)$$

Which ONE of the following statements concerning the above electrolysis is TRUE?

A The reaction is exothermic.

B The reaction is spontaneous.

C chlorine gas is formed at the anode.

D hydrogen gas is formed at the anode. (2)

1.9 Which ONE of the following reactions DOES NOT occur during the Contact Process?

A $S + O_2 \rightarrow SO_2$

B $2\ SO_2 + O_2 \rightarrow 2\ SO_3$

C $SO_3 + H_2O \rightarrow H_2SO_4$

D $SO_3 + H_2SO_4 \rightarrow H_2S_2O_7$ (2)

1.10 Phosphorous rich fertiliser . . .

A improves the quality of fruit and flowers.

B is used to stimulate root growth in plants.

C is produced industrially by the Ostwald process.



D is essential for growing plants with strong stems and healthy green leaves. (2)

[20]

Copyright reserved

QUESTION 2 (Start on a new page.)

The letters A to G in the table below represent seven organic compounds.

A	propanoic acid	**B**	$CH_3CH_2CH_2COCH_2CH_3$
C		**D**	
E	$CH_3CCCH(CH_3)CH_3$	**F**	ethene
G	$CH_3(CH_2)_3CH_3$		

2.1 Using the information in the table, write down the:

2.1.1 IUPAC name of compound **B**. (2)

2.1.2 Name of the FUNCTIONAL GROUP of the homologous series to which compound **A** belongs. (1)

2.1.3 Structural formula of compound **E**. (2)

2.1.4 The type of polymerisation where F is the monomer. (1)

2.1.5 IUPAC name and structural formula of the polymer formed from compound **F**. (3)

2.1.6 Letter that represents a compound that is a chain isomer of 2,2 – dimethylpropane. (1)

2.2 Compound **C**, in the table is the product of the reaction between an alcohol and a carboxylic acid in the presence of concentrated sulphuric acid.

Write down the:

2.2.1 Homologous series to which compound **C** belongs. (1)

2.2.2 IUPAC name of compound **C**. (2)

2.2.3 Structural formula of the carboxylic acid required to prepare compound **C**. (2)

2.2.4 Function of concentrated sulphuric acid in the above reaction. (1)

Copyright reserved

2.3 Compound C, in the table, has isomers.

2.3.1 Define the term *functional isomer.* (2)


2.3.2 Write down the IUPAC name of the straight chain functional isomer of compound **C**. (2)

[20]


QUESTION 3 (Start on a new page.)

Consider the organic compounds represented by the letters **M** and **N** below:

M



N



An investigation is performed to determine which compound; **M** or **N** has a higher vapour pressure.

3.1 Define *vapour pressure.* (2)

3.2 For this investigation write down:

3.2.1 The independent variable. (1)

3.2.2 A controlled variable. (1)

Copyright reserved

3.3 This investigation is carried out at an atmospheric pressure of 100 kPa.

The temperature at which the vapour pressure of compound N equals 100 kPa is 27,80 °C.


3.3.1 At a temperature, T, the vapour pressure of compound **M** equals 100 kPa. Is T, GREATER THAN, EQUAL TO or LESS THAN 27,80 °C? (1)

3.3.2 Refer to structure, intermolecular forces and energy to explain the answer to QUESTION 3.3.1. (3)

3.4 Which compound **M** or **N** has the HIGHER vapour pressure? (1)
[9]

QUESTION 4 (Start on a new page.)

REACTION I and **REACTION II** below are organic reactions in which 2–bromobutane is reacted with solutions of KOH.



Q and **R** are the major organic products formed in **REACTION I** and **REACTION II** respectively.
Q and **R** represent different organic compounds.

4.1 State TWO reaction conditions for:

4.1.1 **REACTION I.** (2)

4.1.2 **REACTION II.** (2)

4.2 What type of reaction (SUBSTITUTION, ADDITION or ELIMINATION), takes place in **REACTION I**? (1)

4.3 Use structural formulae to write down a balanced equation for the reaction that takes place in **REACTION II**. (4)

4.4 Is **R** saturated or unsaturated? Give a reason for the answer. (3)

4.5 In a third reaction, **R** is converted to **Q**. Name the type of addition reaction that takes place during this conversion. (2)
[14]

Copyright reserved

QUESTION 5 (Start on a new page)

Powdered calcium carbonate reacts with hydrochloric acid according to the following balanced equation.

$$CaCO_3(s) + 2HC\ell(aq) \rightarrow CaC\ell_2(aq) + CO_2(g) + H_2O(\ell)$$

The mass of the $CaCO_3$ after every 10 s from the start of the reaction is indicated in the table below:

Time in seconds	0	10	20	30	40	50	60	70	80
Mass of $CaCO_3$ in grams	**X**	74	63	54	46	42	40	40	40

The initial mass of the $CaCO_3$ is **X** g.

5.1 According to the data in the table above, which substance, calcium carbonate or hydrochloric acid is in excess? Give a reason for the answer. (2)

5.2 The average rate of the reaction in the first 30 s is 1,07 $g \cdot s^{-1}$.

5.2.1 Define *reaction rate*, in words. (2)

5.2.2 Calculate the initial mass, **X**, of calcium carbonate. (5)

5.3 What volume of CO_2 is produced between 60 and 80 seconds? (1)

5.4 Use the collision theory to explain why the average rate of reaction between powdered calcium carbonate and the given hydrochloric acid solution decreases with time. (3)

5.5 How will the rate of the above reaction be affected if the initial mass of carbonate was doubled? (Only write down, INCREASES, DECREASES or REMAINS THE SAME) (1)
[14]

Copyright reserved

QUESTION 6 (Start on a new page)

Study the reversible reaction represented by the balanced equation at a fixed temperature below:

$$CH_4(g) + H_2O(g) \rightleftharpoons CO(g) + 3H_2(g) \quad \Delta H = +226 \text{ kJ}$$

Initially, 2,2 moles of $CH_4(g)$ and 1,8 moles of $H_2O(g)$ were mixed in a sealed 2 dm^3 container.

The graph below shows the relationship between the number of moles of CO(g) and time in minutes.



6.1 State Le Chatelier's Principle. (2)

6.2 What information about the reaction can be deduced from the graph after t minutes? (2)

6.3 Calculate the value of the equilibrium constant, Kc, at the fixed temperature used in the reaction. (7)

6.4 Draw a graph (not to scale), on the special answer sheet provided showing the changes in the number of moles of $CH_4(g)$ as the reaction proceeds. Indicate the initial number of moles of $CH_4(g)$ and the number of moles of $CH_4(g)$ at equilibrium, on the graph. (4)

6.5 The initial number of moles of $CH_4(g)$ is now doubled in the same 2 dm^3 container.
How will this change affect the following? (Choose from INCREASES, DECREASES or REMAINS THE SAME)?

6.5.1 The amount of $H_2(g)$ at equilibrium. (1)

6.5.2 Value of the equilibrium constant(Kc). (1)

6.6 Explain the answer to QUESTION 6.5.1 by referring to Le Chatelier's Principle. (3)
[20]

Copyright reserved

QUESTION 7 (Start on a new page.)

7.1 A ***dilute solution of sulphuric acid*** has a concentration 0,012 mol.dm^{-3}

7.1.1 Define an acid according to Arrhenius theory. (2)

7.1.2 Is the above solution of sulphuric acid a strong or weak acid? Explain the answer. (3)

7.2 A learner adds 24 cm^3 of a solution of sulphuric acid of concentration 0,25 mol.dm^{-3} to an unknown volume, X cm^3, of a solution of sodium hydroxide of concentration 0,15 mol.dm^{-3}. The balanced equation for the reaction is:

$$H_2SO_4 + 2NaOH \longrightarrow Na_2SO_4 + 2H_2O$$

The pH of the resulting solution is 0,65.


7.2.1 Calculate the concentration of the sulphuric acid solution after it has reacted with the sodium hydroxide solution. (4)

7.2.2 Calculate the initial volume(X) of the sodium hydroxide solution. Assume that the volume of the solution after the reaction is (X + 24) cm^3. (7)

[16]

QUESTION 8 (Start on a new page.)

A standard galvanic cell is set up using a zinc and an unknown metal, **X**. The graph below shows the change in mass of the metal, **X** whilst the cell is in operation.



8.1 Define an electrolyte. (2)

8.2 Write down the half reaction that takes place at the anode of this cell. (2)

8.3 The initial emf of this cell under standard conditions is 0,63 V.

8.3.1 Write down TWO standard conditions required for the initial emf to be 0,63 V (2)

8.3.2 Identify the metal **X**, by calculating the standard reduction potential of **X**. (5)

8.4 Write down the cell notation for this cell. (3)


8.5 Write down the value of the emf at time **t**, shown on the graph. (1)

[15]

Copyright reserved

QUESTION 9 (Start on a new page.)

An electrolytic cell is set up using two carbon rods, P and Q and a concentrated copper(II) chloride solution as an electrolyte.



When the cell is functioning ONE of the carbon electrodes is coated with a reddish brown layer.

9.1 Write down the energy conversion that takes place in this cell. (1)

9.2 Which electrode, **P** or **Q** is coated with the reddish brown layer? (1)

9.3 Write down a half reaction that takes place at the electrode that is coated with the reddish brown layer. (2)

9.4 The carbon rods in the above cell are **NOW** replaced with copper rods.

9.4.1 Describe the observation that will be made at electrode **Q**. (2)

9.4.2 Refer to relative strengths of reducing agents to explain the observation made at electrode **Q**. (3)

The cell can be used to electroplate a tin medal with a thin layer of copper.

9.4.3 Which electrode, **P** or **Q** must be replaced with the tin medal? (1)
[10]

Copyright reserved

QUESTION 10 (Start on a new page.)

10.1 The unbalanced equations below represent various steps in the Ostwald process.

STEP I $A(g) + O_2(g) \rightarrow B(g) + H_2O(\ell)$

STEP II $B(g) + O_2(g) \rightarrow NO_2(g)$

STEP III $NO_2(g) + H_2O(\ell) \rightarrow C(aq) + NO(g)$

10.1.1 Write down the name of the industrial process that produces gas **A**. (1)


10.1.2 Write down the name or formula of gas **B**. (1)

10.1.3 Name the catalyst that is used in **STEP I**. (1)

The product, **C**, from **STEP III** mixes with reactant A from **STEP I** to produce a fertiliser.

10.1.4 Write a balanced equation to illustrate this reaction. (3)

10.2 Consider the fertiliser in the bag illustrated below:



The only phosphorous containing compound used in the production of this fertiliser is $(NH_4)_3PO_4$

11,95 kg of $(NH_4)_3PO_4$ was used in preparing the fertiliser illustrated above.

Calculate the value represented by the letter **Z**, in the above illustration. (6)

[12]


TOTAL MARKS: 150

Copyright reserved

SPECIAL ANSWER SHEET FOR QUESTION 6.4

NAME: _____ GRADE: _____

DETACH THIS ANSWER SHEET AND SUBMIT WITH YOUR ANSWER BOOKLET



Copyright reserved

DATA FOR PHYSICAL SCIENCES GRADE 12
PAPER 2 (CHEMISTRY)

GEGEWENS VIR FISIESE WETENSKAPPE GRAAD 12
VRAESTEL 2 (CHEMIE)

TABLE 1: PHYSICAL CONSTANTS/*TABEL 1: FISIESE KONSTANTES*

NAME/*NAAM*	SYMBOL/*SIMBOOL*	VALUE/*WAARDE*
Standard pressure *Standaarddruk*	p^θ	$1{,}013 \times 10^5$ Pa
Molar gas volume at STP *Molêre gasvolume by STD*	V_m	22,4 $dm^3 \cdot mol^{-1}$
Standard temperature *Standaardtemperatuur*	T^θ	273 K
Charge on electron *Lading op elektron*	e	$-1{,}6 \times 10^{-19}$ C
Avogadro's constant *Avogadro-konstante*	N_A	$6{,}02 \times 10^{23}$ mol^{-1}

TABLE 2: FORMULAE/*TABEL 2: FORMULES*

$n = \frac{m}{M}$	$n = \frac{N}{N_A}$
$c = \frac{n}{V}$ or/*of* $c = \frac{m}{MV}$	$n = \frac{V}{V_m}$
$\frac{c_a V_a}{c_b V_b} = \frac{n_a}{n_b}$	$pH = -\log[H_3O^+]$
$K_w = [H_3O^+][OH^-] = 1 \times 10^{-14}$ at/*by* 298 K	
$E^\theta_{cell} = E^\theta_{cathode} - E^\theta_{anode}$ / $E^\theta_{sel} = E^\theta_{katode} - E^\theta_{anode}$ or/*of* $E^\theta_{cell} = E^\theta_{reduction} - E^\theta_{oxidation}$ / $E^\theta_{sel} = E^\theta_{reduksie} - E^\theta_{oksidasie}$ or/*of* $E^\theta_{cell} = E^\theta_{oxidising\ agent} - E^\theta_{reducing\ agent}$ / $E^\theta_{sel} = E^\theta_{oksideermiddel} - E^\theta_{reduseermiddel}$	

Copyright reserved

TABLE 3: THE PERIODIC TABLE OF ELEMENTS

KEY/*SLEUTEL*

Atomic number *Atoomgetal*

Symbol *Simbool*

Electronegativity *Elektronegatiwiteit*

Approximate relative atomic mass *Benaderde relatiewe atoommassa*

1 (I)	2 (II)	3	4	5	6	7	8	9	10	11	12	13 (III)	14 (IV)	15 (V)	16 (VI)	17 (VII)	18 (VIII)
1 H 1 (2,1)																	2 He 4
3 Li 7 (1,0)	4 Be 9 (1,5)											5 B 11 (2,0)	6 C 12 (2,5)	7 N 14 (3,0)	8 O 16 (3,5)	9 F 19 (4,0)	10 Ne 20
11 Na 23 (0,9)	12 Mg 24 (1,2)											13 Aℓ 27 (1,5)	14 Si 28 (1,8)	15 P 31 (2,1)	16 S 32 (2,5)	17 Cℓ 35,5 (3,0)	18 Ar 40
19 K 39 (0,8)	20 Ca 40 (1,0)	21 Sc 45 (1,3)	22 Ti 48 (1,5)	23 V 51 (1,6)	24 Cr 52 (1,6)	25 Mn 55 (1,5)	26 Fe 56 (1,8)	27 Co 59 (1,8)	28 Ni 59 (1,8)	29 Cu 63,5 (1,9)	30 Zn 65 (1,6)	31 Ga 70 (1,6)	32 Ge 73 (1,8)	33 As 75 (2,0)	34 Se 79 (2,4)	35 Br 80 (2,8)	36 Kr 84
37 Rb 86 (0,8)	38 Sr 88 (1,0)	39 Y 89 (1,2)	40 Zr 91 (1,4)	41 Nb 92 (1,6)	42 Mo 96 (1,8)	43 Tc (1,9)	44 Ru 101 (2,2)	45 Rh 103 (2,2)	46 Pd 106 (2,2)	47 Ag 108 (1,9)	48 Cd 112 (1,7)	49 In 115 (1,7)	50 Sn 119 (1,8)	51 Sb 122 (1,9)	52 Te 128 (2,1)	53 I 127 (2,5)	54 Xe 131
55 Cs 133 (0,7)	56 Ba 137 (0,9)	57 La 139 (1,0)	72 Hf 179 (1,6)	73 Ta 181 (1,5)	74 W 184 (1,7)	75 Re 186 (1,9)	76 Os 190 (2,2)	77 Ir 192 (2,2)	78 Pt 195 (2,2)	79 Au 197 (2,4)	80 Hg 201 (1,9)	81 Tℓ 204 (1,8)	82 Pb 207 (1,8)	83 Bi 209 (1,9)	84 Po (2,0)	85 At (2,2)	86 Rn
87 Fr (0,7)	88 Ra 226 (0,9)	89 Ac															

58 Ce 140	59 Pr 141	60 Nd 144	61 Pm	62 Sm 150	63 Eu 152	64 Gd 157	65 Tb 159	66 Dy 163	67 Ho 165	68 Er 167	69 Tm 169	70 Yb 173	71 Lu 175
90 Th 232	91 Pa	92 U 238	93 Np	94 Pu	95 Am	96 Cm	97 Bk	98 Cf	99 Es	100 Fm	101 Md	102 No	103 Lr

Copyright reserved

TABLE 4A: STANDARD REDUCTION POTENTIALS
TABEL 4A: STANDAARD-REDUKSIEPOTENSIALE

Increasing oxidising ability/*Toenemende oksiderende vermoë* ↑

Increasing reducing ability/*Toenemende reduserende vermoë* ↓

Half-reactions/*Halfreaksies*	E^{θ} (V)
$F_2(g) + 2e^- \rightleftharpoons 2F^-$	+ 2,87
$Co^{3+} + e^- \rightleftharpoons Co^{2+}$	+ 1,81
$H_2O_2 + 2H^+ + 2e^- \rightleftharpoons 2H_2O$	+1,77
$MnO_4^- + 8H^+ + 5e^- \rightleftharpoons Mn^{2+} + 4H_2O$	+ 1,51
$Cl_2(g) + 2e^- \rightleftharpoons 2Cl^-$	+ 1,36
$Cr_2O_7^{2-} + 14H^+ + 6e^- \rightleftharpoons 2Cr^{3+} + 7H_2O$	+ 1,33
$O_2(g) + 4H^+ + 4e^- \rightleftharpoons 2H_2O$	+ 1,23
$MnO_2 + 4H^+ + 2e^- \rightleftharpoons Mn^{2+} + 2H_2O$	+ 1,23
$Pt^{2+} + 2e^- \rightleftharpoons Pt$	+ 1,20
$Br_2(\ell) + 2e^- \rightleftharpoons 2Br^-$	+ 1,07
$NO_3^- + 4H^+ + 3e^- \rightleftharpoons NO(g) + 2H_2O$	+ 0,96
$Hg^{2+} + 2e^- \rightleftharpoons Hg(\ell)$	+ 0,85
$Ag^+ + e^- \rightleftharpoons Ag$	+ 0,80
$NO_3^- + 2H^+ + e^- \rightleftharpoons NO_2(g) + H_2O$	+ 0,80
$Fe^{3+} + e^- \rightleftharpoons Fe^{2+}$	+ 0,77
$O_2(g) + 2H^+ + 2e^- \rightleftharpoons H_2O_2$	+ 0,68
$I_2 + 2e^- \rightleftharpoons 2I^-$	+ 0,54
$Cu^+ + e^- \rightleftharpoons Cu$	+ 0,52
$SO_2 + 4H^+ + 4e^- \rightleftharpoons S + 2H_2O$	+ 0,45
$2H_2O + O_2 + 4e^- \rightleftharpoons 4OH^-$	+ 0,40
$Cu^{2+} + 2e^- \rightleftharpoons Cu$	+ 0,34
$SO_4^{2-} + 4H^+ + 2e^- \rightleftharpoons SO_2(g) + 2H_2O$	+ 0,17
$Cu^{2+} + e^- \rightleftharpoons Cu^+$	+ 0,16
$Sn^{4+} + 2e^- \rightleftharpoons Sn^{2+}$	+ 0,15
$S + 2H^+ + 2e^- \rightleftharpoons H_2S(g)$	+ 0,14
$2H^+ + 2e^- \rightleftharpoons H_2(g)$	0,00
$Fe^{3+} + 3e^- \rightleftharpoons Fe$	– 0,06
$Pb^{2+} + 2e^- \rightleftharpoons Pb$	– 0,13
$Sn^{2+} + 2e^- \rightleftharpoons Sn$	– 0,14
$Ni^{2+} + 2e^- \rightleftharpoons Ni$	– 0,27
$Co^{2+} + 2e^- \rightleftharpoons Co$	– 0,28
$Cd^{2+} + 2e^- \rightleftharpoons Cd$	– 0,40
$Cr^{3+} + e^- \rightleftharpoons Cr^{2+}$	– 0,41
$Fe^{2+} + 2e^- \rightleftharpoons Fe$	– 0,44
$Cr^{3+} + 3e^- \rightleftharpoons Cr$	– 0,74
$Zn^{2+} + 2e^- \rightleftharpoons Zn$	– 0,76
$2H_2O + 2e^- \rightleftharpoons H_2(g) + 2OH^-$	– 0,83
$Cr^{2+} + 2e^- \rightleftharpoons Cr$	– 0,91
$Mn^{2+} + 2e^- \rightleftharpoons Mn$	– 1,18
$Al^{3+} + 3e^- \rightleftharpoons Al$	– 1,66
$Mg^{2+} + 2e^- \rightleftharpoons Mg$	– 2,36
$Na^+ + e^- \rightleftharpoons Na$	– 2,71
$Ca^{2+} + 2e^- \rightleftharpoons Ca$	– 2,87
$Sr^{2+} + 2e^- \rightleftharpoons Sr$	– 2,89
$Ba^{2+} + 2e^- \rightleftharpoons Ba$	– 2,90
$Cs^+ + e^- \rightleftharpoons Cs$	- 2,92
$K^+ + e^- \rightleftharpoons K$	– 2,93
$Li^+ + e^- \rightleftharpoons Li$	– 3,05

Copyright reserved

TABLE 4B: STANDARD REDUCTION POTENTIALS
TABEL 4B: STANDAARD-REDUKSIEPOTENSIALE

Half-reactions/*Halfreaksies*			E^{θ} (V)
$Li^+ + e^-$	⇌	Li	− 3,05
$K^+ + e^-$	⇌	K	− 2,93
$Cs^+ + e^-$	⇌	Cs	− 2,92
$Ba^{2+} + 2e^-$	⇌	Ba	− 2,90
$Sr^{2+} + 2e^-$	⇌	Sr	− 2,89
$Ca^{2+} + 2e^-$	⇌	Ca	− 2,87
$Na^+ + e^-$	⇌	Na	− 2,71
$Mg^{2+} + 2e^-$	⇌	Mg	− 2,36
$Al^{3+} + 3e^-$	⇌	Al	− 1,66
$Mn^{2+} + 2e^-$	⇌	Mn	− 1,18
$Cr^{2+} + 2e^-$	⇌	Cr	− 0,91
$2H_2O + 2e^-$	⇌	$H_2(g) + 2OH^-$	− 0,83
$Zn^{2+} + 2e^-$	⇌	Zn	− 0,76
$Cr^{3+} + 3e^-$	⇌	Cr	− 0,74
$Fe^{2+} + 2e^-$	⇌	Fe	− 0,44
$Cr^{3+} + e^-$	⇌	Cr^{2+}	− 0,41
$Cd^{2+} + 2e^-$	⇌	Cd	− 0,40
$Co^{2+} + 2e^-$	⇌	Co	− 0,28
$Ni^{2+} + 2e^-$	⇌	Ni	− 0,27
$Sn^{2+} + 2e^-$	⇌	Sn	− 0,14
$Pb^{2+} + 2e^-$	⇌	Pb	− 0,13
$Fe^{3+} + 3e^-$	⇌	Fe	− 0,06
$2H^+ + 2e^-$	⇌	$H_2(g)$	0,00
$S + 2H^+ + 2e^-$	⇌	$H_2S(g)$	+ 0,14
$Sn^{4+} + 2e^-$	⇌	Sn^{2+}	+ 0,15
$Cu^{2+} + e^-$	⇌	Cu^+	+ 0,16
$SO_4^{2-} + 4H^+ + 2e^-$	⇌	$SO_2(g) + 2H_2O$	+ 0,17
$Cu^{2+} + 2e^-$	⇌	Cu	+ 0,34
$2H_2O + O_2 + 4e^-$	⇌	$4OH^-$	+ 0,40
$SO_2 + 4H^+ + 4e^-$	⇌	$S + 2H_2O$	+ 0,45
$Cu^+ + e^-$	⇌	Cu	+ 0,52
$I_2 + 2e^-$	⇌	$2I^-$	+ 0,54
$O_2(g) + 2H^+ + 2e^-$	⇌	H_2O_2	+ 0,68
$Fe^{3+} + e^-$	⇌	Fe^{2+}	+ 0,77
$NO_3^- + 2H^+ + e^-$	⇌	$NO_2(g) + H_2O$	+ 0,80
$Ag^+ + e^-$	⇌	Ag	+ 0,80
$Hg^{2+} + 2e^-$	⇌	$Hg(\ell)$	+ 0,85
$NO_3^- + 4H^+ + 3e^-$	⇌	$NO(g) + 2H_2O$	+ 0,96
$Br_2(\ell) + 2e^-$	⇌	$2Br^-$	+ 1,07
$Pt^{2+} + 2e^-$	⇌	Pt	+ 1,20
$MnO_2 + 4H^+ + 2e^-$	⇌	$Mn^{2+} + 2H_2O$	+ 1,23
$O_2(g) + 4H^+ + 4e^-$	⇌	$2H_2O$	+ 1,23
$Cr_2O_7^{2-} + 14H^+ + 6e^-$	⇌	$2Cr^{3+} + 7H_2O$	+ 1,33
$Cl_2(g) + 2e^-$	⇌	$2Cl^-$	+ 1,36
$MnO_4^- + 8H^+ + 5e^-$	⇌	$Mn^{2+} + 4H_2O$	+ 1,51
$H_2O_2 + 2H^+ + 2e^-$	⇌	$2H_2O$	+1,77
$Co^{3+} + e^-$	⇌	Co^{2+}	+ 1,81
$F_2(g) + 2e^-$	⇌	$2F^-$	+ 2,87

Increasing oxidising ability/*Toenemende oksiderende vermoë* ↓

Increasing reducing ability/*Toenemende reduserende vermoë* ↑

Copyright reserved

education

Department:
Education
PROVINCE OF KWAZULU-NATAL

PHYSICAL SCIENCES P2 (CHEMISTRY)

PREPARATORY EXAMINATION

SEPTEMBER 2019

MEMORANDUM

NATIONAL SENIOR CERTIFICATE

GRADE 12

MARKS : 150

This marking guideline consists of 8 pages.

Copyright Reserved

QUESTION 1

1.1 D ✓✓ (2)

1.2 C✓✓ (2)

1.3 A✓✓ (2)

1.4 C✓✓ (2)

1.5 D✓✓ (2)

1.6 D✓✓ (2)

1.7 D✓✓ (2)

1.8 C✓✓ (2)

1.9 C✓✓ (2)


1.10 B✓✓ (2)
[20]

QUESTION 2

2.1.1 hexan-3-one ✓✓ (2)

2.1.2 carboxyl (group) ✓ (1)

2.1.3




- Whole structure correct: 2/2
- Only functional group correct 1/2
- More than one functional group 0/2

(2)

2.1.4 addition polymerisation✓ (1)

2.1.5



- Whole structure correct ✓✓
- Name ✓


polyethene✓ (3)

2.1.6 G✓ (1)

2.2.1 esters/alkyl alkanoate✓ (1)

2.2.2 ethyl✓ propanoate✓ (2)

2.2.3



- Whole structure correct: 2/2
- Only functional group correct 1/2
- More than one functional group 0/2

(2)

2.2.4 acts as a catalyst/speeds up the reaction. ✓ or
acts as a dehydrating agent. ✓ (1)

2.3.1 Compounds that have the same molecular formula but different functional groups✓✓ (2)

2.3.2 pentanoic acid✓ (2)
[20]

QUESTION 3

3.1 the pressure exerted by a vapour at equilibrium with its liquid in a closed system. ✓✓ (2 or 0) (2)

3.2.1 length of carbon chain/surface area/branching✓ (1)

3.2.2 number of carbon atoms/molecular mass✓ (1)

3.3.1 GREATER THAN✓ (1)

3.3.2 M has a longer carbon chain/greater surface area than N/ M has more sites for intermolecular forces✓
Intermolecular forces between molecules of M are stronger than between molecules of N✓
More energy is required to overcome the intermolecular forces between molecules of M✓ (3)

3.4 N✓ (1)
[9]


QUESTION 4

4.1.1 warm/mild heat✓ dilute KOH✓/warm✓ dilute strong base✓ (2)

4.1.2 hot KOH✓ concentrated ✓Base(KOH) (2)

4.2 substitution✓ (1)

4.3



- ✓ left hand side
- ✓✓ for organic product
- ✓ balancing

(4)

4.4 unsaturated✓
contains a double bond/multiple bond✓ between atoms of carbon✓ (3)

4.5 hydration✓✓ (2)

[14]

QUESTION 5

5.1 calcium carbonate✓ there is some unreacted $CaCO_3$ at the end of the reaction (time 60s) ✓ (2)

5.2.1 **ANY ONE**

- The change in concentration✓ of reactants/products per unit time. ✓
- Rate of change in concentration of reactants or products. ✓✓
- Change in amount/number of moles/volume/mass of reactants/products per (unit) time.
- Amount/number of moles/volume/mass of products formed OR reactants used per (unit) time. (2)

5.2.2 $\text{rate} = -\frac{\text{change in mass of } CaCO_3}{\Delta t}$ ✓

$1{,}07 \overset{\checkmark}{=} -\frac{54 - X}{30 - 0}$ ✓

$= 86{,}10 \text{ g}$ ✓

(if answer is negative minus 1 mark) (5)

Marking criteria

- Equation✓(accept if negative sign is omitted)
- Substitute 54 – X in equation ✓
- Substitute 30 – 0 in equation✓
- Substitute 1,07✓for rate
- Final answer: X = 86,10 g ✓

5.3 0 (cm^3). ✓ (1)

5.4 A decrease in concentration of reactants decreases the number of molecules per unit volume. ✓
Fewer number of collisions per unit time✓
A fewer number of effective collisions occur per unit time/lower frequency of effective collisions. ✓ (3)

5.5 REMAINS THE SAME✓ (1)
[14]

QUESTION 6

6.1 When the equilibrium in a closed system is disturbed, the system will re-instate a new equilibrium by favouring the reaction that will oppose the disturbance. ✓✓ (2 or 0) (2)

6.2 the reaction has reached a state of (dynamic) equilibrium/the rate of the forward reaction is equal to the rate of the reverse reaction. ✓✓(2 or 0) (2)

6.3

Marking criteria:
- Indicating that the number of mols of CO equilibrium is 0,6✓
- Correct mol ratio✓
- Calculating the quantity(mol) at equilibrium of all three substances ✓
- Substitute V = 2 dm^3 in $c = \frac{n}{V}$ to determine concentration at equilibrium of all the substances.✓
- K_c expression✓
- Substitution of concentrations in K_c expression ✓
- Final answer: 0,456 ✓


No K_c expression, correct substitution: Max. $\frac{6}{7}$

Wrong K_c expression : Max. $\frac{4}{7}$

	CH_4	H_2O	CO	H_2	
Initial quantity(mol)	2,2	1,8	0	0	
Change(mol)	-0,6	-0,6	+0,6	+ 1,8	Ratio ✓
Quantity at equilibrium(mol)	1,6	1,2	0,6✓	1,8 ✓	
Equilibrium concentration($mol.dm^{-3}$)	0,8	0,6	0,3	0,9	Divide by 2✓

$$K_c = \frac{[CO]\,[H_2]^3}{[CH_4]\,[H_2O]} \checkmark = \frac{(0,3)\,(0,9)^3}{(0,8)\,(0,6)} \checkmark = 0,456 \checkmark \quad (7)$$

6.4



Graph starts at 2,2✓
Decreases to 1,6✓
Correct shape curve✓
horizontal✓
ignore labels on axes

(4)

6.5.1 INCREASES✓ (1)

6.5.2 REMAINS THE SAME✓ (1)

6.6 An increase in the number of moles of CH_4 increases the concentration of CH_4(reactant).
According to Le Chateliers Principle an <u>increase in the concentration of the reactants</u> ✓ <u>favours the reaction that decreases the concentration of the reactants</u>✓
In this case the <u>forward reaction is favoured</u>✓

(3)
[20]

QUESTION 7

7.1.1 An acid is a substance that produces hydrogen ions(H^+)/hydronium ions(H_3O^+)✓ when it dissolves in water. ✓ (2)

7.1.2 strong✓
it ionises completely in water✓ ✓ (ACCEPT: dissociates) (3)

7.2.1

$pH = - \log [H_3O^+]$ ✓

0,65✓ $= - \log [H_3O^+]$

$\therefore [H_3O^+] = 0,224$ mol.dm^{-3}

$c((H_2SO_4) = \frac{1}{2}c(H_3O^+)$ ✓
$= \frac{1}{2}(0,224)$
$= 0,112$ mol.dm^{-3}✓

- Formula pH = - log $[H_3O^+]$ ✓
- Substitute 0,65 for pH✓
- $c((H_2SO_4) = \frac{1}{2}c(H_3O^+)$✓
- $c((H_2SO_4) = 0,112$ mol.dm^{-3}✓

(4)

7.2.2 POSITIVE MARKING FROM QUESTION 7.2.1: concentration of H_2SO_4

Marking guidelines/*Nasienriglyne:*

- Formulae: $c = \frac{n}{V}$ /n = cV/ ✓
- Calculate initial number of moles of H_2SO_4✓
- Calculate number of moles of H_2SO_4 that reacted✓
- Calculate number of moles of H_2SO_4 in excess✓
- Calculate number of moles of NaOH that reacted✓
- Ratio of NaOH to H_2SO_4✓
- Final answer cm^3 or dm^3✓

$n(H_2SO_4)initial = cV$✓
$= (0,25)(0,024)$ ✓
$= 6 \times 10^{-3}$ mols

$n(H_2SO_4)excess = cV$
$= (0,112)\frac{(X + 24)}{1000}$ ✓

$n(H_2SO_4)reacted = 6 \times 10^{-3} - \frac{(0,112)(X + 24)}{1000}$ ✓

$n(NaOH)reacted = cV$
$\frac{0,15(X)}{1000}$ ✓

$n(NaOH)reacted = 2\ (n(H_2SO_4)reacted)$✓

$\frac{0,15(X)}{1000} = 2\left(6 \times 10^{-3} - \frac{(0,112)(X + 24)}{1000}\right)$
$X = 17,71\ cm^3$✓ $0,01771\ dm^3$

(7)
[16]

QUESTION 8

8.1 a solution/liquid/dissolved substance✓ that conducts electricity through the movement of ions.✓ (2)

8.2 $\underline{Zn(s) \rightarrow Zn^{2+}(aq) + 2e^-}$

Notes

- $Zn^{2+} + 2e^- \leftarrow Zn$ $(2/2)$ $Zn^{2+} + 2e^- \rightleftharpoons Zn$ $(0/2)$
  $Zn \rightleftharpoons Zn^{2+} + 2e^-$ $(1/2)$ $Zn^{2+} + 2e \rightarrow Zn$ $(0/2)$
- Ignore if charge on electron is omitted.
- If a charge of an ion is omitted e.g. $Zn \rightarrow Zn^2 + 2e^-$ Max.: $1/2$

(2)

8.3.1 Temperature of 25 ºC/298K✓
Concentration of the electrolytes equals 1 $mol.dm^{-3}$. ✓ (2)

8.3.2 $E^{\Theta}_{cell} = E^{\Theta}_{cathode} - E^{\Theta}_{anode}$ ✓
0,63✓ $= E^{\Theta}_{cathode} - (-0,76)$ ✓
$E^{\Theta}_{cathode} = -0,13$ V ✓
X is lead(Pb) ✓

Notes
- Accept any other correct formula from the data sheet.
- Any other formula using unconventional abbreviations, e.g. $E°_{cell} = E°_{OA} - E°_{RA}$ followed by correct substitutions:
  $E°_{sel} = E°_{OM} - E°_{RM}$ *Max/:* $^{4}/_{5}$

(5)

8.4 $Zn(s)/Zn^{2+}(aq)$ ✓//✓$Pb^{2+}(aq)/Pb(s)$ ✓
$Zn(s)/Zn^{2+}(1\ mol.dm^{-3})$ ✓//✓$Pb^{2+}(1\ mol.dm^{-3})/Pb(s)$
Accept $Zn/Zn^{2+}//Pb^{2+}/Pb$ (3)

8.5 0(V) ✓ (1)

[15]

QUESTION 9

9.1 from electrical energy to chemical energy✓ (1)

9.2 P✓ (1)

9.3 $Cu^{2+} + 2e^{-} \rightarrow Cu$

Notes
- $Cu^{2+} + 2e^{-} \leftarrow Cu$ ($^{0}/_{2}$) $Cu^{2+} + 2e^{-} \rightleftharpoons Cu$ ($^{1}/_{2}$)
  $Cu \rightleftharpoons Cu^{2+} + 2e^{-}$ ($^{0}/_{2}$) $Cu^{2+} + 2e \rightarrow Cu$ ($^{2}/_{2}$)
- Ignore if charge on electron is omitted.
- If a charge of an ion is omitted e.g. $Cu \rightarrow Cu^{2} + 2e^{-}$ Max.: $^{1}/_{2}$

(2)

9.4.1 Q will break down/become eroded/surface becomes rough and eroded✓✓
ACCEPT Q will be oxidised. ✓✓ (2)

9.4.2 Cu/electrode Q is a stronger reducing agent✓ than the $C\ell^{-}$ ions✓.
Cu/Q will be oxidised/loses electrons✓ resulting in the electrode becoming eroded
OR
The $C\ell^{-}$ ion is a weaker reducing agent✓ than Cu(Q) ✓and will therefore not be oxidised. ✓ (3)

9.4.3 P✓ (1)

[10]

QUESTION 10

10.1.1 Haber process✓ (1)
10.1.2 nitric oxide✓ NO✓ (1)
10.1.3 platinum✓ (1)
10.1.4 $HNO_3 + NH_3 \rightarrow (NH_4)NO_3$ (3)

Notes:
- Reactants ✓ Products ✓ Balancing ✓
- Ignore double arrows.
- Marking rule 6.3.10.

10.2
Mass P = 31✓/149✓ x11,95✓ = 2,486 kg
2/8 x Z/100 x 20✓ = 2,486✓
Z = 49,72%✓

(6)

TOTAL MARKS: 150

Copyright Reserved